



Take the Guesswork Out of Oral Appliance Therapy

- ✓ Settings can be precisely transferred to a custom appliance
- ✓ Fitted by any trained healthcare professional
- Avoids components that can be swallowed, lost, or dropped
- ✓ Same silicone retention material used by dentists to take impressions
- Comprehensive training videos at: www.youtube.com/user/sleepusa









Accommodates full range of arch widths with 3 vertical sizes





2237 Faraday Ave, Suite 100 Carlsbad, CA 92008 P 760.720.0099 | F 760.476.3620

and catalyst. Mix amounts of base consistent color. Roll in the bottom tray with hands and place thoroughly to a Measure equal



and distribute the of the bottom tray evenly to the edges 2. Press down retention material



3. Instruct the prepared bottom tray is now bite down firmly for Apnea Guard and patient to insert the 1.5 minutes. The



Maximum work time - 1.5 minutes from start of mixing

6. With the Apnea

8. Instruct the

Minimum set time - 1.5 minutes

their lower jaw patient to protrude

set 1 minute after 4. Repeat steps mixing begins material will start to remember the 1-2 for the top tray



prepare the <u>top</u> tray bites down firmly ensure the patient 5. Repeat Step 3 and for 1.5 minutes to



setting number on a permanent marker the tray handle with 7. Write the neutral



setting number on 9. Write the max setting number the tray handle below the neutral

on the look-up table the neutral and maximum settings to determine the 10. Locate 70% (optimal) setting



70% setting to fix the trays at the 1 1. Insert the lock



to be worn during Guard is now ready

14. The Apnea

cover into place 13. Slide the lock



regularly and to stop the Apnea Guard using after 29 days patient to rinse 15. Instruct the